

7.) $m = 10 \text{ kg}$ $F_1(x=1) = 20 \text{ N}$ $F_2 = -b\dot{x}$ $v_0 = 0$

$F_1 = -Dx$ $D = 20 \frac{\text{N}}{\text{m}}$ $\boxed{\vec{a} = \frac{\vec{F}_e}{m}}$ $x(t=3T) = \frac{x(0)}{10}$

$T = ?$

$m\ddot{x} = -Dx - b\dot{x}$

$$\omega_0^2 = \frac{D}{m} \quad \frac{b}{m} = 2\alpha \quad \ddot{x} + \frac{b}{m}\dot{x} + \frac{D}{m}x = 0$$

$$\ddot{x} + 2\alpha\dot{x} + \omega_0^2 x = 0$$

$$x(t) = A e^{-\alpha t} \cos(\omega t + \delta) \quad \omega = \sqrt{\omega_0^2 - \alpha^2}$$

$$x(0) = A \cos \delta$$

$$x(3T) = A e^{-3\alpha T} \cos(3T\omega + \delta) = A e^{-3\alpha T} \cos \delta$$

$$\frac{x(0)}{x(3T)} = 10 = \frac{A \cos \delta}{A e^{-3\alpha T} \cos \delta} = e^{3\alpha T} \rightarrow \ln 10 = 3\alpha T$$

$$\frac{\ln 10}{3T} = \alpha$$

$$\omega^2 = \omega_0^2 - \alpha^2 \quad \omega = \frac{2\pi}{T}$$

$$\omega_0^2 - \alpha^2 = \frac{4\pi^2}{T^2}$$

$$\omega_0^2 - \left(\frac{\ln 10}{3T}\right)^2 = \frac{4\pi^2}{T^2}$$

↓

T