


5.)

1. oldal

 $H(r) = ?$ ábrázolni

Ampère-féle gerj. tv.



$$\oint_G \vec{H} \cdot d\vec{s} = \sum I_i = \int_A \vec{j} \cdot d\vec{A}$$

 $r \leq r_0$: $\vec{j}$ homogén és kifelé $j_1 = \frac{I}{r_0^2 \pi}$

$$\oint_G \vec{H} \cdot d\vec{s} = \int_A \vec{j} \cdot d\vec{A}$$


$$H \cdot 2r\pi = \frac{I}{r_0^2 \pi} r^2 \pi$$

$$\Downarrow$$

 $H = \dots$
 $r_0 < r \leq r_1$ $\vec{j} = 0$
ha $r > r_0$ 

$$\oint_G \vec{H} \cdot d\vec{s} = \int_A \vec{j} \cdot d\vec{A}$$

$$H \cdot 2r\pi = \frac{I}{r_0^2 \pi} r_0^2 \pi = I \rightarrow H = \dots$$

 $r_1 < r \leq r_2$ $\vec{j}$ homogén és befelé

$$A_2 = r_2^2 \pi - r_1^2 \pi$$

$$j_2 = \frac{I}{r_2^2 \pi - r_1^2 \pi} = \frac{I}{(r_2^2 - r_1^2) \pi}$$

$$\oint_G \vec{H} \cdot d\vec{s} = \int_A \vec{j} \cdot d\vec{A}$$

$$H \cdot 2r\pi = \frac{I}{(r_2^2 - r_1^2) \pi} (r^2 - r_1^2) \pi = \dots$$